

DTFT

$$x[n] = \frac{1}{2\pi} \int_{-\pi}^{\pi} X(e^{j\omega}) e^{j\omega n} d\omega$$

$$X(e^{j\omega}) = \sum_{n=-\infty}^{\infty} x[n] e^{-j\omega n}$$

Symmetry Properties

| | |
|--------------------------|--|
| $x[n]$ | $X(e^{j\omega})$ |
| $x^*[n]$ | $X^*(e^{-j\omega})$ |
| $x^*[-n]$ | $X^*(e^{j\omega})$ |
| $x[-n] = x^*[n]$ | $X(e^{j\omega}) = X^*(e^{j\omega})$ (Real) |
| $x[-n] = -x^*[n]$ | $X^*(e^{j\omega}) = -X(e^{j\omega})$ (Imaginary) |
| $x[n] = x_e[n] + x_o[n]$ | $Re\{X(e^{j\omega})\} + jIm\{X(e^{j\omega})\}$ |
| | Conjugate Symmetric |
| | Real part is even |
| Real x[n] | Imaginary part is odd |
| | Magnitude is even |
| | Phase is odd |

Theorems

| | |
|---|---|
| $x[n]$ | $X(e^{j\omega})$ |
| $\alpha x[n] + \beta y[n]$ | $\alpha X(e^{j\omega}) + \beta Y(e^{j\omega})$ |
| $x[n - n_d]$ | $e^{-j\omega n_d} X(e^{j\omega})$ |
| $x[-n]$ | $X(e^{-j\omega})$ |
| $nx[n]$ | $j \frac{dX(e^{j\omega})}{d\omega}$ |
| $x[n] * y[n]$ | $X(e^{j\omega}) Y(e^{j\omega})$ |
| $x[n]y[n]$ | $\frac{1}{2\pi} \int_{-\pi}^{\pi} X(e^{j\theta}) Y(e^{j(\omega-\theta)}) d\theta$ |
| $\sum_{n=-\infty}^{\infty} x[n] ^2 = \frac{1}{2\pi} \int_{-\pi}^{\pi} X(e^{j\omega}) ^2 d\omega$ | |
| $\sum_{n=-\infty}^{\infty} x[n]y^*[n] = \frac{1}{2\pi} \int_{-\pi}^{\pi} X(e^{j\omega}) Y^*(e^{j\omega}) d\omega$ | |
| Converges if $\sum_{n=-\infty}^{\infty} x[n] < \infty$ | |

Transform Pairs

| Signal | Fourier Transform |
|---|--|
| $\delta[n]$ | 1 |
| $\delta[n - n_0]$ | $e^{-j\omega n_0}$ |
| 1 | $\sum_{k=-\infty}^{\infty} 2\pi \delta(\omega + 2\pi k)$ |
| $a^n u[n] (\alpha < 1)$ | $\frac{1}{1 - \alpha e^{-j\omega}}$ |
| $u[n]$ | $\frac{1}{1 - e^{-j\omega}} + \sum_{k=-\infty}^{\infty} \pi \delta(\omega + 2\pi k)$ |
| $(n+1)a^n u[n]$ | $\frac{1}{(1 - \alpha e^{-j\omega})^2}$ |
| $\frac{\sin \omega_c n}{\pi n}$ | $\begin{cases} 1 & \omega < \omega_c, \\ 0 & \omega_c < \omega \leq \pi \end{cases}$ |
| $\begin{cases} 1 & 0 \leq n \leq M, \\ 0 & \text{else} \end{cases}$ | $\frac{\sin(\omega(M+1)/2)}{\sin[\omega]/2} e^{-j\omega M/2}$ |
| $e^{j\omega_0 n}$ | $\sum_{k=-\infty}^{\infty} 2\pi \delta(\omega - \omega_0 + 2\pi k)$ |
| $\cos(\omega_0 n + \phi)$ | $\sum_{k=-\infty}^{\infty} \pi e^{j\phi} \delta(\omega - \omega_0 + 2\pi k) + \pi e^{-j\phi} \delta(\omega + \omega_0 + 2\pi k)$ |

Z - Transform

$$X(z) = \sum_{n=-\infty}^{\infty} x[n]z^{-n}$$

Theorems

| Sequence | Transform | ROC |
|---------------------|----------------------------------|----------------|
| $ax_1[n] + bx_2[n]$ | $aX_1(z) + bX_2(z)$ | $R_1 \cap R_2$ |
| $x[n - n_0]$ | $z^{-n_0}X(z)$ | R |
| $z_0^n x[n]$ | $X\left(\frac{z}{z_0}\right)$ | $ z_0 R$ |
| $nx[n]$ | $-z \frac{dX(z)}{dz}$ | R |
| $x^*[n]$ | $X^*(z^*)$ | R |
| $Re\{x[n]\}$ | $\frac{1}{2} [X(z) + X^*(z^*)]$ | Contains R |
| $Im\{x[n]\}$ | $\frac{1}{2j} [X(z) - X^*(z^*)]$ | Contains R |
| $x^*[-n]$ | $X^*\left(\frac{1}{z^*}\right)$ | $\frac{1}{R}$ |
| $x_1[n] * x_2[n]$ | $X_1(z)X_2(z)$ | $R_1 \cap R_2$ |

Transforms

| Sequence | Transform | ROC |
|--|--|-----------------------------------|
| $\delta[n]$ | 1 | \mathbb{C} |
| $u[n]$ | $\frac{1}{1-z^{-1}}$ | $ z > 1$ |
| $-u[-n - 1]$ | $\frac{1}{1-z^{-1}}$ | $ z < 1$ |
| $\delta[n - m]$ | z^{-m} | \mathbb{C} except 0 or ∞ |
| $a^n u[n]$ | $\frac{1}{1-az^{-1}}$ | $ z > a $ |
| $-a^n u[-n - 1]$ | $\frac{1}{1-az^{-1}}$ | $ z < a $ |
| $na^n u[n]$ | $\frac{az^{-1}}{(1-az^{-1})^2}$ | $ z > a $ |
| $-na^n u[-n - 1]$ | $\frac{az^{-1}}{(1-az^{-1})^2}$ | $ z < a $ |
| $r^n \cos(\omega_0 n)u[n]$ | $\frac{1-r\cos(\omega_0)z^{-1}}{1-2r\cos(\omega_0)z^{-1}+r^2z^{-2}}$ | $ z > r$ |
| $r^n \sin(\omega_0 n)u[n]$ | $\frac{r\sin(\omega_0)z^{-1}}{1-2r\cos(\omega_0)z^{-1}+r^2z^{-2}}$ | $ z > r$ |
| $\left\{ \begin{array}{l} a^n \quad 0 \leq n \leq N-1 \\ 0 \end{array} \right\}$ | $\frac{1-a^N z^{-N}}{1-az^{-1}}$ | $ z > 0$ |

ROC properties

1. ROC is either $|z| \geq r$, $|z| \leq r$, or $r_1 \leq |z| \leq r_2$
2. Fourier transform exists iff ROC includes unit circle
3. ROC contains no poles
4. If $x[n]$ is finite, ROC is entire plane
5. If $x[n]$ is right sided, ROC extends from outermost pole to ∞
6. If $x[n]$ is left sided, ROC extends from innermost pole to 0
7. If $x[n]$ is two sided, then ROC is an annulus
8. ROC is connected region
9. $\sum_{n=-\infty}^{\infty} |x[n]| \implies$ ROC includes unit circle

For LTI Systems

10. Stable \implies Causal if and only if right-sided ROC
11. Causal \implies Stable if and only if poles inside unit circle

Discrete Fourier Series

$$\tilde{x}[n] = \frac{1}{N} \sum_k \tilde{X}[k] e^{j \frac{2\pi}{N} kn}$$

$$\tilde{X}[n] = \sum_n \tilde{x}[n] e^{-j \frac{2\pi}{N} kn}$$

Properties

| Periodic sequence $\tilde{x}[n]$ | Periodic coefficients $\tilde{X}[k]$ |
|---|---|
| $a\tilde{x}_1[n] + b\tilde{x}_2[n]$ | $a\tilde{X}_1[k] + b\tilde{X}_2[k]$ |
| $\tilde{X}[n]$ | $N\tilde{x}[-k]$ |
| $\tilde{x}[n - m]$ | $e^{-j \frac{2\pi}{N} km} \tilde{X}[k]$ |
| $e^{j \frac{2\pi}{N} mn} \tilde{x}[n]$ | $\tilde{X}[k - m]$ |
| $\sum_{m=0}^{N-1} \tilde{x}_1[m] \tilde{x}_2[n - m]$ | $\tilde{X}_1[k] \tilde{X}_2[k]$ |
| $\tilde{x}^*[n]$ | $\tilde{X}^*[-k]$ |
| $\tilde{x}^*[-n]$ | $\tilde{X}^*[k]$ |
| $Re\{\tilde{x}[n]\}$ | $\tilde{X}_e[k] = \frac{1}{2} (\tilde{X}[k] + \tilde{X}^*[-k])$ |
| $jIm\{\tilde{x}[n]\}$ | $\tilde{X}_o[k] = \frac{1}{2} (\tilde{X}[k] - \tilde{X}^*[-k])$ |
| $\tilde{x}_e[n] = \frac{1}{2} (\tilde{x}[n] + \tilde{x}^*[-n])$ | $Re\{\tilde{X}[k]\}$ |
| $\tilde{x}_o[n] = \frac{1}{2} (\tilde{x}[n] - \tilde{x}^*[-n])$ | $jIm\{\tilde{X}[k]\}$ |
| | Conjugate Symmetric |
| | Real Part Even |
| | Imaginary Part Odd |
| | Magnitude is even |
| | Phase is odd |
| Real $\tilde{x}[n]$ | |

Discrete Fourier Transform

$$x[n] = \frac{1}{N} \sum_{k=0}^{N-1} X[k] e^{j \frac{2\pi}{N} kn}$$

$$X[k] = \sum_{n=0}^{N-1} x[n] e^{-j \frac{2\pi}{N} kn}$$

Properties

| Finite Length sequence $x[n]$ | N Point DFT $X[k]$ |
|---|---|
| $ax_1[n] + bx_2[n]$ | $aX_1[k] + bX_2[k]$ |
| $X[n]$ | $Nx[((-k))_N]$ |
| $x[((n - m))_N]$ | $e^{-j \frac{2\pi}{N} km} X[k]$ |
| $e^{j \frac{2\pi}{N} mn} x[n]$ | $X[((k - m))_N]$ |
| $\sum_{m=0}^{N-1} x_1[m] x_2[((n - m))_N]$ | $X_1[k] X_2[k]$ |
| $x^*[n]$ | $X^*[((-k))_N]$ |
| $x^*[((-n))_N]$ | $X^*[k]$ |
| $Re\{x[n]\}$ | $X_e[k] = \frac{1}{2} (X[k] + X^*[-k])$ |
| $jIm\{x[n]\}$ | $X_o[k] = \frac{1}{2} (X[k] - X^*[-k])$ |
| $x_e[n] = \frac{1}{2} (x[n] + x^*[((-n))_N])$ | $Re\{X[k]\}$ |
| $x_o[n] = \frac{1}{2} (x[n] - x^*[((-n))_N])$ | $jIm\{X[k]\}$ |
| | Conjugate Symmetric |
| | Real Part Even |
| | Imaginary Part Odd |
| | Magnitude is even |
| | Phase is odd |
| Real $x[n]$ | |

$$\sum_{n=0}^{N-1} |x[n]|^2 = \frac{1}{N} \sum_{k=0}^{N-1} |X[k]|^2$$

$$x[n] = \frac{1}{N} DFT\{X^*[k]\}^*$$